

Package Leaflet: Information for the User**E M Pharma One A Day Hayfever Relief
10 mg Tablets
Loratadine**

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist have told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in the leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse.

What is in this leaflet:

- 1. What this medicine is and what it is used for**
- 2. What you need to know before you take this medicine**
- 3. How to take this medicine**
- 4. Possible side effects**
- 5. How to store this medicine**
- 6. Contents of the pack and other information**

1 What this medicine is and what it is used for

- E M Pharma One A Day Hayfever Relief 10 mg Tablets contains the active substance loratadine
- loratadine belongs to a group of drugs called antihistamines which help to relieve the symptoms of some allergies
- your medicine is used to relieve symptoms such as sneezing, runny nose and burning, itchy eyes due to hayfever and other allergies. This medicine can also be taken to relieve symptoms due to urticaria (hives) such as rash, itching and redness.

2 What you need to know before you take this medicine**Do NOT take this medicine if you:**

- are allergic to loratadine or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions**Talk to your doctor or pharmacist before you start taking this medicine if you:**

- have severe liver problems (see section 3, 'How to take this medicine').

Other precautions you should take

This product may prevent a response to allergens in skin allergy testing; therefore you should stop taking this medicine at least 2 days before any such testing.

Other medicines and E M Pharma One A Day Hayfever Relief 10 mg Tablets

Talk to your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Pregnancy and breast-feeding

Do not take this medicine if you are pregnant or breast-feeding.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

This medicine may make you feel drowsy. Do not drive or operate any tools or machines if affected.

This medicine contains lactose

Patients who are intolerant to lactose should note that this medicine contain a small amount of lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3 How to take this medicine

If your doctor has prescribed this medicine, always take this medicine exactly as your doctor has told you. Otherwise follow the instructions below. You should check with your doctor or pharmacist if you are not sure.

The tablets should be swallowed, preferably with a drink of water.

The usual dose is:

Adults and Children over 12 years of age
One 10 mg tablet once daily.

Children aged 2 - 12 years of age are dosed by weight:

Body weight more than 30 kg: 10 mg once daily (one tablet once daily).

This medicine are not suitable for children with a body weight of less than 30 kg. For children aged 2 years and over but weighing 30 kg or less, a syrup form of this medicine may be more suitable.

Children under 2 years of age

This medicine is not recommended for children under 2 years of age.

Patients with severe liver problems

Adults and children over 2 years of age
Bodyweight more than 30kg: 10 mg (one tablet) every other day.

Top of page cut-off to middle of registration mark: 21mm.

If you take more of this medicine than you should

If you (or someone else) swallow a lot of the tablets all together, or if you think a child has swallowed any of the tablets, contact your nearest hospital casualty department or your doctor immediately. Overdose may cause sleepiness, fast heart rate and headache. Please take this leaflet, any remaining tablets and the container with you to the hospital or doctor so that they know which tablets were consumed.

If you forget to take this medicine

If you forget to take a dose, take one as soon as you remember, unless it is nearly time to take the next one. Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this **medicine**, ask your doctor or pharmacist.

4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If the following happens, stop taking this medicine and tell your doctor immediately or go to the casualty department at your nearest hospital:

- a severe allergic reaction (swelling of the face, lips, mouth or throat which may cause difficulty swallowing or breathing).

This is a very serious but very rare side effect. You may need urgent medical attention or hospitalisation.

The following side effects have been reported at the approximate frequencies shown:

Children aged 2 to 12 years:

Common (may affect up to 1 in 10 people):

- headache
- nervousness
- tiredness (fatigue).

Adults and children over 12 years:

Common (may affect up to 1 in 10 people):

- drowsiness (somnolence)
- headache
- increased appetite
- difficulty sleeping (insomnia).

Very rare (may affect up to 1 in 10,000 people):

- dizziness
- convulsions
- fast or uneven heartbeat (tachycardia, palpitations)
- feeling sick (nausea), dry mouth, inflammation of the stomach, which may lead to stomach pain, being sick, or loss of

- appetite (gastritis)
- abnormal liver function (shown by blood test results)
- rash, hair loss (alopecia)
- tiredness (fatigue).

Not known (cannot be estimated from available data)

- weight increased

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

5 How to store this medicine

Keep this medicine out of the sight and reach of children. This medicinal product does not require any special storage conditions. Do not transfer the tablets to another container. Do not use this medicine after the expiry date that is stated on the outer packaging. The expiry date refers to the last day of that month. Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

Contents of the pack and other information**What this medicine contain:**

- the active ingredient is loratadine 10 mg
- the other ingredients are **lactose**, maize starch, pregelatinised starch and magnesium stearate.

What this medicine look like and contents of the pack:

- white, oval tablets, scored on one side and plain on the other, debossed "L" and "10" on each side of the scoreline
- the product is available in packs of 5, 7, 10, 14, 15, 20, 28 and 30 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation holder and company responsible for manufacture: TEVA UK Limited, Eastbourne, BN22 9AG.

This leaflet was last revised: December 2017

PL 00289/0371

87914-B EAS1980a
160 x 210