

Aspirin 300 MG Tablets 16's 'Pull Out' leaflet LABEL (Seven Panel) V7 (St Albans)

Tablets Item Code AVL020/10

NEW LABEL-Leaflet Layout

Read attached leaflet for full instructions before use

Usage: For the relief of mild to moderate pain including headache, migraine, neuralgia

(nerve pain), bothache, sore throat and period pains.

For the symptomatic relief of colds and flu, rheumatic pain, sprains and strains, muscular aches, fibrositis, sciatica, lumbago, joint swelling and stiffness.

How to take: Check the bottle seal is not broken before first use. If it is do not take the tablets. For oral use, The tablets should be swallowed with a drink of water.

Dosage: Adults, the elderly and children over 16 years: 1 to 3 tablets to be swallowed with water every 4 hours, or as directed by your doctor. Not more than 4 doses in any 24 hours.

16 years: Take 1 to 3 tablets with water, every 4 hours. Do not take more

Ingredients:
Each Tablet contains active: aspirin 300mg.
Also contains inactive: lactose, potato starch and talc,

3. HOW TO TAKE THE TABLETS

SAVED AS FINAL ARTWORK 30.05.18 **ILLUSTRATOR CS6** FONTS TURNED TO OUTLINE PRINT FILES SUPPLIED AS PRESS QUALITY PDF & Illustrator CS6.ai Please check where applicable any

BARCODES/PHARMACODES/ITEM CODES

[SPECIAL ADDRESSES]

V I C T O R D E S MANUFACTURER PRODUCT Aspirin 300 MG Tablets 16's 'Pull Out' leaflet LABEL PL NUMBER 08977/0002, 08977/0018, 08977/0022 and 08977/0023 A/W VERSION Seventh draft (St.A) [Special addresses] DIMENSIONS Multi-panel profile see ARTWORK for full detail DATE tel: 01275 342028 • email: mail@victor-design.com

Aspirin 300mg tablets are available in blister packs of 2, 4, 8,

12, 16, 24, 32, 48, 50, 96 or 100 tablets and they are also available in container packs of 8, 16, 24, 25, 32, 50,100, 250, 500, 1000 or 5000 tablets (Not all pack sizes are marketed).

Date of text revision: June 2017_ _ _ _ _ _

Page 7

with client prior to final print Important warning:

Do not give this medicine to children under the age of 16 unless your doctor tells you. This is because there is a possible association between Aspirin and Reye's Syndrome when given to children. Reye's Syndrome is a very rare disease, which can be fatal.

Take special care and tell your doctor if you:

Are intolerant to some sugars as these tablets contain lactose

Have asthma or have ever had it in the past

Have dehydration

Metoclopramide [for sickness and digestive disorders]

Corticosteroids e.g. hydrocortisone (which can be used arthritis, asthma and inflammatiory conditions)

Anti-inflammatory medicines or steroids for pain and inflammation e.g. ibuprofer

High blood pressure medicines e.g. diuretic/water tablets and a first material to the past and the past a A lower dose is recommended for the elderly.

Do not give to children under 16 years of age unless on the advice of a doctor,
If symptoms persist for more than 3 days consult your doctor. rheumatic pain, sciatica, lumbago, muscle aches, fibrositis, Patient Information Leaflet for Aspirin 300 mg Tablets Blood thinning medicines sucin as nepatin, trainant or incommalone
Corticosteroids e.g., hydrocortisone (which can be used for arthritis, asthma and inflammatory conditions)
Anti-inflammatory medicines or steroids for pain and inflammation e.g., ibuprofen
High blood pressure medicines e.g. diuretic/water tablets, Angio-I receptor antagonists or ACE inhibitors
Epilepsy medicines e.g. Phenytoin, Sodium valproate
Methotrexate (for psoriasis, arthritis or tumours)
Antacids for indigestion
Probenecid or sulphinpyrazone for gout
Anti-depressants e.g. oftalopram Read this leaflet carefully before taking these tablets.

It does not contain all the information about your medicine that you may need to know, so ask your doctor or pharmacist if you have any questions.

It is leaflet only applies to Aspirin 300 mg Tablets. 2. CHECK BEFORE YOU TAKE ASPIRIN 300 ma Do not take these tablets if you:

Are allergic to Aspirin, NSAIDs which includes those when in attacks of asthma, angioedema, uticaria or rhinitis have been precipitated by aspirin or any other NSAID or any of the other ingredients listed in section 6

Hawe a stomach ulcer or have had problems with ulcers in DO NOT EXCEED THE STATED DOSE TABLETS Do not take if you are allergic to aspirin or any other anti-inflammate pain killers or any of the other ingredients, if you have or ever had stomach ulcer, or have or ever had gout or blood clotting disorders Consult your doctor before taking aspirin if you are pregnant or breastleeding, you suffer from asthma or are taking other medication. Have dehydration
 Have been told you are deficient in glucose-6-phosphate 1. WHAT THESE TABLETS DO WHAT THESE TABLETS DO
These tablets contain Aspirin which is a type of painkiller. It is
used to relieve pain, swelling and high temperature. It can be
used to treat headaches, migraine, neuralgia, toothache, sore
throat, period pains and for the relief of sprains, strains, Suffer from gout or had in the past
 Suffer from gout or had in the past
 Are in the last 3 months of pregnancy or are breast feeding. lehydrogenase I ydurg doctor or pharmacist if you are already taking y of the following medicines, as their effectiveness may influenced by aspirin if they are taken at the same time.

Page 1 Page 3 Page 4 106 mm 106 mm 106 mm than 4 dösés in any 24 höurs.
A lower dose is recommended for the elderly.
When taking this medicine, it is important to remember the following:
• Do not give to children aged under 16 years unless told to by your doctor. 4. POSSIBLE SIDE EFFECTS These other effects are less serious. 5. STORING YOUR MEDICINE Manufacturer Before using aspirin inform your doctor about the medicines These other effects are less serious.
If they bother you talk to a pharmacist:
Feeling sick, being sick, changes to the stomach lining (swelling or ulcers)
Increased bleeding time when you cut yourself, bleeding under the skin which may look like bruising
Nose bleeds (if a nose bleed is severe or lasts for a long time, talk to a doctor straight away)
Hearing problems (e.g. ringing in the ears) feeling dizzy (vertigo), feeling confused before using aspirin inform your octor about the medicines you are taking. If you are using aspirin regularly seek advice of your doctor before taking any other medicine (including other medicines that you have bought).

Pregnancy and breast-feeding
Ask your doctor or pharmacist for advice before taking this medicine. Aspirin should not be taken in the last three months of pregnancy. Do not take this medicine if you are breastfeeding. Aspar Pharmaceuticals Ltd,
29 - 30 Capitol Way, Colindale, London, NW9 0EQ.
PL 08977/0002, PL 08977/0018, PL 08977/0022 and PL 08977/0023 Keep your medicine in a safe place where children cannot see or reach it. Most people will not have problems, but some may get some.
If you get any of these serious side effects, stop taking the
tablets. See a doctor at once:

You are sick and it contains blood or dark particles that look Do not use this medicine after the expiry date printed on the MA Holder holder and manufacturer is by your doctor
Do not exceed the stated dose
Seek medical attention IMMED YOU are sick and it Contains broud or usin particles that born like coffee grounds
 Pass blood in your stools or pass black tarry stools
 Stomach problems including pain, indigestion or heartburn
 Alergic reactions such as skin rash, swelling of the face, neck or throat, worsening of asthma, difficulty in breathing
 Bleeding on the brain which may cause a severe headache Aspar Pharmaceuticals Ltd. Acrewood Way, St Albans, AL4 0JY, United Kingdom. tion IMMEDIATELY if you accidentally 6. MORE ABOUT YOUR MEDICINE

Page 6

for the missed dose

If symptoms persist for more than 3 days consult your doctor

120 mm ASPAID Aspirin 300 mg Tablets For oral use. The tablets should be swallowed with a drink of Do not take if you are alleroic to aspirin or any other Read attached leaflet for full instructions before use. anti-inflammatory pain killers or any of the other ingredients, if you have or ever had a stomach ulcer, or have or ever had a stomach ulcer, or have or ever had gout or blood clotting disorders. Consult your doctor before taking aspirin if you are pregnant or breastfeeding, you suffer 1 to 3 tablets to be swallowed with water every 4 hours, or as directed by your doctor. Not more than 4 doses in any 24 hours. Usage:
For the relief of mild to moderate pain including headache, migraine, neuralgia (nerve pain), toothache, sore throat and period pains. For the symptomatic relief of colds and flu, rheumatic pain, sprains and strains, muscular aches, fibrositis, sciatica, lumbago, joint swelling and stiffness. Store in the original package.

KEEP OUT OF THE SIGHT AND REACH OF CHILDREN. ersist for more than 3 days consult your doctor, PL Holder:
Aspar Pharmaceuticals Ltd, St Albans, AL4 0JY
PL 08977/0002
GSL DO NOT EXCEED THE STATED DOSE.

Page 5

If you miss a dose, do not take a double dose to make up.


Page 2 (base label)

Braille text reads as follows:

aspirin # 3 0 0 m q tablets

denotes unique number symbol


Note: dies comply with Marburg Medium cell dimensions:



Please Check all Barcodes, Pharmacodes and item codes PRIOR to final Printing, thank you



PLEASE REMOVE THIS CUTTER GUIDE BEFORE PRINTING



Technical & Non Printing Items Cutter Guides




Aspirin 300 MG Tablets 16's 'Pull Out' leaflet LABEL (Seven Panel)



NEW LABEL-Leaflet Layout


BRAILLE ONLY



Print Colours



(K) black 100%



This LABEL - Leaflet was originaly created on 17.03.14, please Check all Barcodes, Pharmacodes and item codes PRIOR to final Printing, thank you






Aspirin 300 MG Tablets 100's 'Pull Out' leaflet LABEL (Seven Panel) V7

100 Tablets Item Code AVL028/8

NEW LABEL-Leaflet Layout




Braille text reads as follows:

Aspirin 300 mg Tablets


Read attached leaflet for full instructions before use.
Ingredients:
Each Tablet contains active: aspirin 300mg.
Also contains inactive: aspirin 300mg

Page 2 (base label)

PLEASE REMOVE THIS CUTTER GUIDE BEFORE PRINTING



Note: dies comply with Marburg Medium cell dimensions:



This LABEL - Leaflet was originaly created on 17.03.14, please Check all Barcodes, Pharmacodes and item codes PRIOR to final Printing, thank you

C T O R D E S

Multi-panel profile see ARTWORK for full detail

Aspirin 300 MG Tablets 100's 'Pull Out' leaflet LABEL

08977/0002, 08977/0018, 08977/0022 and 08977/0023

Technical & Non Printing Items

Print Colours

P485

(K) black 100%

MANUFACTURER PRODUCT

PL NUMBER

INKS USED

tel: 01275 342028 • email: mail@victor-design.com

A/W VERSION
DIMENSIONS






Aspirin 300 MG Tablets 100's 'Pull Out' leaflet LABEL (Seven Panel)



NEW LABEL-Leaflet Layout

BRAILLE ONLY




Cutter Guides

Colours

Adhesive_Deadener Braille

Print Colours



This LABEL - Leaflet was originaly created on 17.03.14, please Check all Barcodes, Pharmacodes and item codes PRIOR to final Printing, thank you

