



Aspirin DISPERSIBLE 75MG Tablets 100's 'Pull Out' leaflet LABEL (Seven Panel) V15 (D'cant)

100 Tablets

 ITEM CODE DA013/10 (TBA 19.11.18)

"do not eat desiccant"

V I C T O R D E S I G N	
MANUFACTURER	Aspar Pharmaceuticals Limited
PRODUCT	Aspirin DISPERSIBLE 75MG Tablets 100's 'Pull Out' leaflet LABEL
PL NUMBER	08977/0007
A/W VERSION	Fifteenth draft (D'cant)
DIMENSIONS	Multi-panel profile see ARTWORK for full detail
INKS USED	Black - Pantone 368 C Green - Cyan
DATE	19.11.18
tel: 01275 342028 • email: mail@victor-design.com	

NEW LABEL-Leaflet Layout

Technical Et Non Printing Items

Cutter Guides

Colours

Adhesive	3	Braille	4
Deadener			

BLACK	
SPOT 1	PANTONE 368 C
SPOT 2	CYAN 100%

107 mm | 106 mm | 106 mm

32 mm

Page 1 | Page 3 | Page 4

106 mm | 106 mm | 106 mm

32 mm

Page 5 | Page 6 | Page 7

120 mm


32 mm

Peel and Lift

Page 2 (base label)


Braille text reads as follows:

d i s p e r s i b l e
a s p i r i n t a b l e t s
7 5 m g



denotes unique number symbol

Note: dies comply with Marburg Medium cell dimensions:



This LABEL - Leaflet was originally created on 17.03.14, please Check all Barcodes, Pharmacodes and item codes PRIOR to final Printing, thank you

PLEASE REMOVE THIS CUTTER GUIDE BEFORE PRINTING





Aspirin DISPERSIBLE 75MG Tablets 100's 'Pull Out' leaflet LABEL (Seven Panel)

100 Tablets

BRaille ONLY


NEW LABEL-Leaflet Layout

Technical Et Non Printing Items
Cutter Guides

Colours

Adhesive_Deadener	Braille
3	4

BLACK
SPOT 1 PANTONE 368 C



Page 1

Page 3


Page 4



Page 5

Page 6

Page 7




Page 2 (base label)


Braille text reads as follows:

d i s p e r s i b l e
a s p i r i n t a b l e t s
7 5 m g

denotes unique number symbol



Note: dies comply with Marburg Medium cell dimensions:



(Figures are in mm)

PLEASE REMOVE THIS CUTTER GUIDE BEFORE PRINTING

This LABEL - Leaflet was originally created on 17.03.14, please Check all Barcodes, Pharmacodes and item codes PRIOR to final Printing, thank you





Aspirin DISPERSIBLE 75MG Tablets 1000's 'Pull Out' leaflet LABEL (Seven Panel) V9 (d'cant)

1000 Tablets

ITEM CODE DA011/8 (TBA 19.11.18)

NEW LABEL-Leaflet Layout

"do not eat desiccant"

V I C T O R D E S I G N	
MANUFACTURER	Aspar Pharmaceuticals Limited
PRODUCT	Aspirin DISPERSIBLE 75MG Tablets 1000's 'Pull Out' leaflet LABEL
PL NUMBER	08977/0007
A/W VERSION	Ninth draft (d'cant)
DIMENSIONS	Multi-panel profile see ARTWORK for full detail
INKS USED	Black - Pantone 368 C Green - Pantone 287 C - Cyan
DATE	19.11.18
tel: 01275 342028 • email: mail@victor-design.com	

Technical Et Non Printing Items

Cutter Guides

Colours

Adhesive	Deadener	Braille
3		4

BLACK	
SPOT 1	PANTONE 368 C
SPOT 2	CYAN 100%
SPOT 3	PANTONE 287 C

107 mm 106 mm 106 mm

32 mm

Read attached leaflet for full instructions before use.

DISPERSIBLE ASPIRIN TABLETS 75 mg

Helps prevent blood clots in people who have had a stroke, heart attack or by-pass surgery

1000 tablets

The usual dose, for long term use, is 1 or 2 tablets once daily. In some circumstances a higher dose may be appropriate, especially in the short term and up to 4 tablets a day may be used on the advice of a doctor.

Do not give to children aged under 16 years, unless on the advice of a doctor.

Warnings (DO NOT EXCEED THE STATED DOSE)

Do not take these tablets if you:

• Are allergic to Aspirin, NSAIDs which includes those when in attacks of asthma, angioedema, urticaria or rhinitis have been precipitated by aspirin or any other NSAID or any of the other ingredients listed in section 6

• Suffer from gout or had in the past

• Have a stomach ulcer or have had problems with ulcers in the past

• Suffer from haemophilia or any other bleeding condition

2. CHECK BEFORE YOU TAKE

Before starting long term use of Aspirin you must consult your doctor who will discuss the benefits and risks of this therapy with you.

Do not take these tablets if you:

• Are taking blood thinning (anti-coagulant) medication

• Are in the last 3 months of pregnancy or are breast feeding.

Important warning:

Do not give this medicine to children under the age of 16 unless your doctor tells you. This is because there is a possible association between Aspirin and Reye's Syndrome when given to children. Reye's syndrome is a very rare disease, which can be fatal.

Take special care and tell your doctor if you:

• Have liver or kidney disease

• Have asthma or have ever had it in the past

• Are suffering from dehydration (feeling constantly thirsty)

• Are intolerant to some sugars as these tablets contain lactose

• Have been told you are deficient in glucose-6-phosphate dehydrogenase

Tell your doctor or pharmacist if you are already taking any of the following medicines, as their effectiveness may be influenced by aspirin if they are taken at the same time.

• Metoclopramide (for sickness and digestive disorders)

• Blood thinning medicines such as Heparin, Warfarin or nicoumalone

• Corticosteroids e.g. hydrocortisone (which can be used for arthritis, asthma and inflammatory conditions)

• Anti-inflammatory medicines or steroids for pain and inflammation e.g. ibuprofen

• High blood pressure medicines e.g diuretic/water tablets, Angio-II receptor antagonists or ACE inhibitors

• Epilepsy medicines e.g. Phenytoin, Sodium valproate

Page 1 Page 3 Page 4

106 mm 106 mm 106 mm

32 mm

3. HOW TO TAKE THE TABLETS

Do not take these tablets without discussing it with your doctor first.

Dosage in adults, the elderly and children over the age of 16 years:

The usual dosage is 1 to 2 tablets dissolved in water, once a day. In some people a higher dose of up to 4 tablets a day may be prescribed.

Do not take this many tablets unless told to by your doctor. When taking this medicine, it is important to remember the following:

• The tablets must be dissolved in a glass of water before taking

4. POSSIBLE SIDE EFFECTS

Most people will not have problems, but some may get some. If you get any of these serious side effects, stop taking the tablets. See a doctor at once:

• You are sick and it contains blood or dark particles that look like coffee grounds

• Pass blood in your stools or pass black tarry stools

• Stomach problems including pain, indigestion or heartburn

• Allergic reactions such as skin rash, swelling of the face, neck or throat, worsening of asthma, difficulty in breathing

• Bleeding on the brain which may cause a severe headache or stroke

These other effects are less serious. If they bother you talk to a pharmacist:

• Feeling sick, being sick, changes to the stomach lining (swelling or ulcers)

• Increased bleeding time when you cut yourself, bleeding under the skin which may look like bruising

• Nose bleeds (if a nose bleed is severe or lasts for a long time, talk to a doctor straight away)

5. STORING YOUR MEDICINE

Keep your medicine in a safe place where children cannot see or reach it.

Do not store above 25°C.

Keep the product in the original container in order to protect from moisture.

Do not use this medicine after the expiry date printed on the packaging.

6. MORE ABOUT YOUR MEDICINE

The active ingredient in these tablets is 75mg aspirin. The inactive ingredients are starch, sodium saccharin, lactose, citric acid, calcium carbonate, talc and sodium lauryl sulphate.

The product licence holder and manufacturer is Aspar Pharmaceuticals Ltd, 29-30 Capitol Way, Colindale, NW9 0EQ, UK. PL 08977/0007

Manufacturer: Aspar Pharmaceuticals Ltd, Albany House, Acrewood Way, St Albans, AL4 0JY, United Kingdom.

Dispensible Aspirin 75mg tablets are available in blister packs of 24, 28 and 100 tablets and they are also available in container packs of 25, 28, 32, 50, 100 or 1000 tablets (Not all pack sizes are marketed). Date of text revision: November 2018.

Page 5 Page 6 Page 7

120 mm

32 mm

Peel and Lift

Dispensible Aspirin Tablets 75 mg

Read attached leaflet for full instructions before use.

Ingredients:

Each tablet contains: Aspirin 75mg. Also contains lactose and sodium.

Usage: an antiplatelet agent which helps to prevent blood clots in people who have had a stroke, heart attack or by-pass surgery.

How to take:

Check the bottle seal is not broken before first use. If it is do not take the tablets. The tablets should be dispersed in water before taking by mouth.

Dosage:

The advice of a doctor should be sought before commencing therapy for the first time.

The usual dose, for long term use, is 1 or 2 tablets once daily. In some circumstances a higher dose may be appropriate, especially in the short term and up to 4 tablets a day may be used on the advice of a doctor.

Do not give to children aged under 16 years, unless on the advice of a doctor.

Warnings

DO NOT EXCEED THE STATED DOSE.

Do not take these tablets if you:

• Are allergic to Aspirin, NSAIDs which includes those when in attacks of asthma, angioedema, urticaria or rhinitis have been precipitated by aspirin or any other NSAID or any of the other ingredients listed in section 6

• Suffer from gout or had in the past

• Have a stomach ulcer or have had problems with ulcers in the past

• Suffer from haemophilia or any other bleeding condition

2. CHECK BEFORE YOU TAKE

Before starting long term use of Aspirin you must consult your doctor who will discuss the benefits and risks of this therapy with you.

Do not take these tablets if you:

• Are taking blood thinning (anti-coagulant) medication

• Are in the last 3 months of pregnancy or are breast feeding.

Important warning:

Do not give this medicine to children under the age of 16 unless your doctor tells you. This is because there is a possible association between Aspirin and Reye's Syndrome when given to children. Reye's syndrome is a very rare disease, which can be fatal.

Take special care and tell your doctor if you:

• Have liver or kidney disease

• Have asthma or have ever had it in the past

• Are suffering from dehydration (feeling constantly thirsty)

• Are intolerant to some sugars as these tablets contain lactose

• Have been told you are deficient in glucose-6-phosphate dehydrogenase

Tell your doctor or pharmacist if you are already taking any of the following medicines, as their effectiveness may be influenced by aspirin if they are taken at the same time.

• Metoclopramide (for sickness and digestive disorders)

• Blood thinning medicines such as Heparin, Warfarin or nicoumalone

• Corticosteroids e.g. hydrocortisone (which can be used for arthritis, asthma and inflammatory conditions)

• Anti-inflammatory medicines or steroids for pain and inflammation e.g. ibuprofen

• High blood pressure medicines e.g diuretic/water tablets, Angio-II receptor antagonists or ACE inhibitors


• Epilepsy medicines e.g. Phenytoin, Sodium valproate

Page 2 (base label)


Braille text reads as follows:

dispersible
aspirin tablets
75 mg

denotes unique number symbol



Note: dies comply with Marburg Medium cell dimensions:



(Figures are in mm)

PLEASE REMOVE THIS CUTTER GUIDE BEFORE PRINTING





Aspirin DISPERSIBLE 75MG Tablets 1000's 'Pull Out' leaflet LABEL (Seven Panel)

1000 Tablets

BRaille ONLY

NEW LABEL-Leaflet Layout


PLEASE NOTE: 11.05.15 Braille should be placed as far right as possible, so as not to impede 'BATCH & EXP' overprinting by Aspar

Technical & Non Printing Items

Cutter Guides

Colours


Adhesive_Deadener	3	Braille	4



Page 1

Page 3


Page 4



Page 5

Page 6

Page 7




Page 2 (base label)


Braille text reads as follows:

d i s p e r s i b l e
a s p i r i n t a b l e t s
7 5 m g

denotes unique number symbol



Note: dies comply with Marburg Medium cell dimensions:



(Figures are in mm)

PLEASE REMOVE THIS CUTTER GUIDE BEFORE PRINTING

