



MercuryPharma

Patient Information Leaflet

Phorpain® Gel Maximum Strength

Ibuprofen



↑
pharmacode read direction

Read all of this leaflet carefully before you start taking this medicine.

Keep this leaflet. You may need to read it again. If you have any further questions, ask your doctor or pharmacist. This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours. If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What is Phorpain Gel Maximum Strength and what it is used for
2. Before you use Phorpain Gel Maximum Strength
3. How to use Phorpain Gel Maximum Strength
4. Possible side effects
5. How to store Phorpain Gel Maximum Strength
6. Further information

1. WHAT IS PHORPAIN GEL MAXIMUM STRENGTH AND WHAT IT IS USED FOR

Phorpain Gel Maximum Strength contains ibuprofen and belongs to a group of medicines called non-steroid anti-inflammatory drugs (NSAIDs). These medicines reduce pain and inflammation, and bring down a high temperature.

Phorpain Gel Maximum Strength is used to treat a number of painful conditions affecting the joints and muscles such as back pain, sprains, strains and other sport injuries. It is also used for the relief of pain caused by mild to moderate arthritic conditions and nerve pain (neuralgia).

2. BEFORE YOU USE PHORPAIN GEL MAXIMUM STRENGTH

DO NOT use Phorpain Gel Maximum Strength:

- If you are allergic to ibuprofen, aspirin or similar medicines (e.g NSAIDs) or any of the other ingredients in this gel (listed at the end of this leaflet)
- If you are asthmatic, or suffer from rhinitis (allergic runny nose) or urticaria (hives)
- On broken, damaged, infected or diseased skin.

Take special care with Phorpain Gel Maximum Strength

- Protect treated areas from direct sunlight to avoid any sensitivity reaction, e.g. a rash
- You have had an ulcer or some other problem affecting your stomach, or intestines in the past

- You have asthma or wheezing attacks (or if you have had asthma in the past)
- You have any kidney problems
- If you develop a rash after using the gel stop using it any further
- If you suffer from bronchial asthma or any allergic disease.
- Do not smoke or go near naked flames - risk of severe burns. Fabric (clothing, bedding, dressings etc) that has been in contact with this product burns more easily and is a serious fire hazard. Washing clothing and bedding may reduce product build-up but not totally remove it.

If any of the above apply to you, only use this product on advice from your doctor or pharmacist.

PHORPAIN GEL IS FOR EXTERNAL USE ONLY

Taking / Using other medicines:

Can you take Phorpain Gel with other medicines?

Please inform your doctor or pharmacist if you are taking, or have recently taken any other medicine, even those not prescribed by a doctor.

The effect of this medicine may affect or be affected by taking the following medicines at the same time:

- Medicines to lower your blood pressure (e.g. atenolol)
- Medicines used to thin the blood (e.g. warfarin)
- Aspirin or other NSAIDs (including other forms of Ibuprofen), used for pain and inflammation.

Please tell your doctor or pharmacist if you are taking any of the above, or have recently taken any other medicine - even those not prescribed.

Pregnancy and breast feeding

You should not use Phorpain Gel Maximum Strength if you are pregnant or breast-feeding.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Phorpain Gel Maximum Strength should not affect the ability to drive or use machines.

3. HOW TO USE PHORPAIN GEL MAXIMUM STRENGTH

Phorpain Gel Maximum Strength is designed for topical (on the skin) application only. Never take the gel by mouth.

Always use Phorpain Gel Maximum Strength exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

Check the seal is intact before first use (invert cap to break seal).

Adults: Squeeze 2 to 5cm (i.e. 0.8 to 2 inches) of gel from the tube on affected area.

Massage until absorbed. This dose should not be repeated more frequently than every four hours and no more than four times a day in any 24 hour period. The amount of gel squeezed would be equivalent to 50 to 125mg of ibuprofen.

Phorpain Gel Maximum Strength should only be used on healthy, unbroken skin. Do not use it on or near cuts or grazes or under dressings such as plasters. Also, do not use it on the genital area.

Do not let any gel come in contact with your eyes. If it does, rinse your eyes with cold water and consult your doctor. Hands should be washed after applying Phorpain Gel Maximum Strength, unless they are the site of treatment.

If the condition does not improve after two weeks use, or becomes worse at any time, speak to your doctor or pharmacist.

Children: Phorpain Gel Maximum Strength is not recommended for use in children under 12 years.

If the gel is accidentally swallowed:

If you accidentally swallow any Phorpain Gel Maximum Strength, rinse out your mouth thoroughly and contact your doctor, or nearest hospital, as soon as possible. If swallowed, the gel may cause an upset stomach.

If you forget to use your Phorpain Gel Maximum Strength:

If you miss a dose, just carry on with the next dose as normal. Do not apply a double dose.

4. POSSIBLE SIDE EFFECTS

Like all medicines Phorpain Gel can cause side effects, although not everybody gets them.

All medicines can cause allergic reactions although serious allergic reactions are rare. Any sudden wheeziness, difficulty in breathing, swelling of the eyelids, face or lips, rash or itching (especially affecting your whole body) should be reported to a doctor immediately.

Other side effects include:

- Itching or reddening of the skin
- Abdominal Pain (pains in your stomach) or other abnormal stomach symptoms
- A burning feeling
- Sore or weeping spots
- Kidney problems (particularly in people who have a history of kidney disease).

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard
By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE PHORPAIN GEL MAXIMUM STRENGTH

Keep this medicine out of the sight and reach of children.

Do not use Phorpain Gel Maximum Strength after the expiry date which is stated on the tube. The expiry date refers to the last day of that month. It should be stored in a cool, dry place, below 25°C. Keep the tube tightly closed.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Remember:

This medicine is for you. Never give this medicine to someone else, it could harm them, even if their symptoms seem the same as yours.

6. FURTHER INFORMATION

What Phorpain Gel Maximum Strength contains:

The active substance is ibuprofen. Other ingredients are hydroxyethylcellulose, sodium hydroxide, benzyl alcohol, isopropyl alcohol, purified water.

What Phorpain Gel Maximum Strength looks like and contents of the pack:

Phorpain Gel Maximum Strength is supplied in aluminium tubes containing 100g gel.

Marketing Authorisation Holder

Mercury Pharma Group Ltd.,
Capital House, 85 King William Street,
London EC4N 7BL, UK

Manufacturer:

Mercury Pharmaceuticals Ltd.,
Capital House, 85 King William Street,
London EC4N 7BL, UK

Alternate Manufacturer:

Farmasierra Manufacturing S.L.
Carretera De Irun KM 26,200 28700,
SAN Sebastian De Los Reyes, Madrid, Spain.

This leaflet was last revised in August 2019.

Phorpain is the registered trade mark of Mercury Pharma Group Ltd.

© Mercury Pharma Group Ltd. 2000

LF-118849-01
17006/9