

56 mm

154 mm

Rennie Liquid Heartburn Relief forms a protective layer on top of the stomach contents to prevent stomach acid escaping into the food pipe which causes pain and discomfort.

Dosage: For oral use. Shake well before use.

Adults and children over 12 years: Take 10-20 ml (two to four 5 ml spoonfuls) after meals and at bedtime, up to four times a day.

Children 6 to 12 years: Take 5-10 ml (one to two 5 ml spoonfuls) after meals and at bedtime, up to four times a day.

Children under 6 years: Not recommended.

See leaflet for further information.

This product is sugar free.

You can take this product if you are pregnant or breast-feeding.

LIQUID HEARTBURN RELIEF

oral suspension

Sodium Alginate,
Sodium Bicarbonate,
Calcium Carbonate

150 ml

MINT flavour

Fast, soothing relief from heartburn & indigestion

Each 5 ml of this medicine contains:

133.5 mg Sodium Bicarbonate, 250 mg Sodium Alginate, 80 mg Calcium Carbonate. It also contains ethyl parahydroxybenzoate (E214), propyl parahydroxybenzoate (E216), butyl parahydroxybenzoate.

Keep out of the sight and reach of children.

Do not store above 25°C.
Do not refrigerate.

MA Holder and Manufacturer:
Pinewood Laboratories Ltd.,
Ballymacarbry, Clonmel,
Co. Tipperary, Ireland.
PL 04917/0027

Distributed by:
Bayer plc, RG2 6AD.

231103324W

B/N:

EXP:

Peel Here

5 010605 300936

Liquid Heartburn Relief

Oral Suspension

Peppermint Flavour

Sodium Bicarbonate, Sodium Alginate, Calcium Carbonate

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you. Always take this medicine exactly as described in the leaflet or as your doctor or pharmacist has told you.

What Rennie® Liquid Heartburn Relief is and what it is used for

Rennie Liquid Heartburn Relief belongs to a group of medicines called 'reflux suppressants' which form a protective layer on top of the stomach contents to prevent stomach acid escaping from the stomach where it works into the food pipe causing pain and discomfort.

This liquid is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy.

Before using Rennie® Liquid Heartburn Relief

Do not take this product if:

- You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).

Take special care before treatment with Rennie® Liquid Heartburn Relief:

This medicine contains small amounts of sodium (6 mmol per 10 ml). If you have been advised to follow a low sodium (salt) diet ask your doctor before taking this product. This product contains ethyl parahydroxybenzoate (E214), propyl parahydroxybenzoate (E216), and butyl parahydroxybenzoate which may cause an allergic reaction (possibly delayed).

Rennie Liquid Heartburn Relief may be used during pregnancy and breast-feeding.

Taking other medicines

Do not take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and beta blockers (for heart conditions), penicillamine (for rheumatoid arthritis), glucocorticoid (for inflammatory and autoimmune disorders), neuroleptics

(for mental illness), thyroid hormones, chloroquine (for malaria), estramustine (for prostate cancer) and bisphosphonates (for osteoporosis).

How to take Rennie® Liquid Heartburn Relief

To be taken by mouth, after meals and at bedtime. Shake bottle well before use.

Adults and children over 12 years: Take 10-20 ml (two to four 5 ml spoonfuls) after meals and at bedtime, up to four times a day.

Children 6 to 12 years: Take 5-10 ml (one to two 5 ml spoonfuls) after meals and at bedtime, up to four times a day.

Children under 6 years: Not recommended.

If you take too much of this product you may feel bloated. It is unlikely to cause any harm, but please consult your doctor or pharmacist. If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.

After taking Rennie® Liquid Heartburn Relief

If symptoms persist after 7 days consult your doctor.

Possible side effects

Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, difficulty in breathing, dizziness or swelling of the face, lips, tongue or throat.

Reporting of side effects:

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

How to store Rennie® Liquid Heartburn Relief

Keep all medicines out of the sight and reach of children.

Do not use after the expiry date shown on the pack (EXP month/year). Do not store above 25°C. Do not refrigerate.

Further information

Each 5 ml of this medicine contains 133.5 mg Sodium Bicarbonate, 250 mg Sodium Alginate, 80 mg Calcium Carbonate. It also contains carbomer, sodium hydroxide, saccharin sodium, ethyl parahydroxybenzoate (E214), propyl parahydroxybenzoate (E216), butyl parahydroxybenzoate, isopropyl alcohol, peppermint oil and purified water.





MA Holder and Manufacturer

Pinewood Laboratories Ltd., Ballymacarbry, Clonmel, Co. Tipperary, Ireland.

Revision date: December 2018

Bayer





56 mm

154 mm

Rennie Liquid Heartburn Relief forms a protective layer on top of the stomach contents to prevent stomach acid escaping into the food pipe which causes pain and discomfort.

Dosage: For oral use. Shake well before use.

Adults and children over 12 years: Take 10-20 ml (two to four 5 ml spoonfuls) after meals and at bedtime, up to four times a day.

Children 6 to 12 years: Take 5-10 ml (one to two 5 ml spoonfuls) after meals and at bedtime, up to four times a day.

Children under 6 years: Not recommended.

See leaflet for further information.

This product is sugar free.

You can take this product if you are pregnant or breast-feeding.

Rennie

LIQUID HEARTBURN RELIEF

oral suspension

Sodium Alginate, Sodium Bicarbonate, Calcium Carbonate

250 ml

MINT flavour

Fast, soothing relief from heartburn & indigestion

BAYER

Each 5 ml of this medicine contains:

133.5 mg Sodium Bicarbonate, 250 mg Sodium Alginate, 80 mg Calcium Carbonate. It also contains ethyl parahydroxybenzoate (E214), propyl parahydroxybenzoate (E216), butyl parahydroxybenzoate.

Keep out of the sight and reach of children.

Do not store above 25°C.

Do not refrigerate.

MA Holder and Manufacturer:

Pinewood Laboratories Ltd., Ballymacabry, Clonmel, Co. Tipperary, Ireland.

PL 04917/0027

Distributed by:

Bayer plc, RG2 6AD.

Peel Here

5 010605 300967

B/N:

EXP:

2310525PW

Rennie

Liquid Heartburn Relief

Oral Suspension

Peppermint Flavour

Sodium Bicarbonate, Sodium Alginate, Calcium Carbonate

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you. Always take this medicine exactly as described in the leaflet or as your doctor or pharmacist has told you.

What Rennie® Liquid Heartburn Relief is and what it is used for

Rennie Liquid Heartburn Relief belongs to a group of medicines called 'reflux suppressants' which form a protective layer on top of the stomach contents to prevent stomach acid escaping from the stomach where it works into the food pipe causing pain and discomfort.

This liquid is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy.

Before using Rennie® Liquid Heartburn Relief

Do not take this product if:

- You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).

Take special care before treatment with Rennie® Liquid Heartburn Relief:

This medicine contains small amounts of sodium (6 mmol per 10 ml). If you have been advised to follow a low sodium (salt) diet ask your doctor before taking this product. This product contains ethyl parahydroxybenzoate (E214), propyl parahydroxybenzoate (E216), and butyl parahydroxybenzoate which may cause an allergic reaction (possibly delayed).

Rennie Liquid Heartburn Relief may be used during pregnancy and breast-feeding.

Taking other medicines

Do **not** take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and beta blockers (for heart conditions), penicillamine (for rheumatoid arthritis), glucocorticoid (for inflammatory and autoimmune disorders), neuroleptics

(for mental illness), thyroid hormones, chloroquine (for malaria), estramustine (for prostate cancer) and bisphosphonates (for osteoporosis).

How to take Rennie® Liquid Heartburn Relief

To be taken by mouth, after meals and at bedtime. Shake bottle well before use.

Adults and children over 12 years: Take 10-20 ml (two to four 5 ml spoonfuls) after meals and at bedtime, up to four times a day.

Children 6 to 12 years: Take 5-10 ml (one to two 5 ml spoonfuls) after meals and at bedtime, up to four times a day.

Children under 6 years: Not recommended.

If you take too much of this product you may feel bloated. It is unlikely to cause any harm, but please consult your doctor or pharmacist. If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.

After taking Rennie® Liquid Heartburn Relief

If symptoms persist after 7 days consult your doctor.

Possible side effects

Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, difficulty in breathing, dizziness or swelling of the face, lips, tongue or throat.

Reporting of side effects:

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

How to store Rennie® Liquid Heartburn Relief

Keep all medicines out of the sight and reach of children.

Do not use after the expiry date shown on the pack (EXP month/year). Do not store above 25°C. Do not refrigerate.

Further information

Each 5 ml of this medicine contains 133.5 mg Sodium Bicarbonate, 250 mg Sodium Alginate, 80 mg Calcium Carbonate. It also contains carbomer, sodium hydroxide, saccharin sodium, ethyl parahydroxybenzoate (E214), propyl parahydroxybenzoate (E216), butyl parahydroxybenzoate, isopropyl alcohol, peppermint oil and purified water.

MA Holder and Manufacturer


Pinewood Laboratories Ltd., Ballymacabry, Clonmel, Co. Tipperary, Ireland.

Revision date: December 2018

Bayer

MOCK-UP!

not for production



BRAILLE CONVERSION – English

A large rectangular area filled with numerous small green dots arranged in a complex, non-random pattern. The dots are distributed across the entire width and height of the section, creating a dense field of points.

Rennie Liquid Heartburn Relief Oral Suspension