

FRADOR

CHLORBUTOL 1.0% w/v

- Gives prompt relief from the pain associated with mouth ulcers and gum sores and on-going protection from saliva, bacteria and food whilst healing proceeds.
 - Frador should be applied 4 times a day after meals and before going to bed.
- In the event of missing an application, apply **as needed**.

DIRECTIONS:

Stand bottle in tray provided. Dry Ulcer. Using applicator, apply Frador directly to mouth ulcer. There may be a temporary tingling sensation. Keep mouth open for several seconds to allow Frador to dry. Application is complete. If healing does not start within a few days always consult your doctor.

CAUTION:

As with all medicines - KEEP OUT OF THE SIGHT AND REACH OF CHILDREN. **For adult use only.** Avoid use in pregnancy. Keep away from naked flame. Retain carton for instructions. Not to be taken internally. In rare occurrences users can experience a local reaction to some or all of the ingredients. If excess irritation occurs during or after application, stop use. If condition persists, contact your pharmacist or doctor. There have been no reports of ill-effects from overdosage. Store at room temperature.

REPORTING SIDE EFFECTS:

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

Licence holder: Fenton Pharmaceuticals, Fenton House, 4 Hampstead Gate, 1a Frognal, London, NW3 6AL
Contract Manufacturer: Thorpe Laboratories LN12 1NB.

FRADOR

Expiry Date:
Batch No:

Active ingredients: Chlorbutol B.P. 1.0% w/v
Excipients: Menthol B.P. 0.1% w/v, Prep Syrax E.P.C. 2.9% w/v,
Balsamic Benzoin 15.0% w/v, in Alcohol Solution to 100% Carmoisine.

G.S.L.

FRADOR
For mouth ulcers

For mouth ulcers and gum sores

- Immediate effective relief
- Soothes pain & aids healing
- Hygienic application

3.5 ml

CHLORBUTOL 1.0% w/v

FRADOR



PL 24784/0001

FRADOR