

PATIENT LEAFLET: INFORMATION FOR THE USER

NUMARK HAYFEVER RELIEF FOR ADULTS 50 MICROGRAMS / DOSE NASAL SPRAY

Beclometasone Dipropionate

The name of your medicine is Numark Hayfever Relief for Adults 50 micrograms / dose Nasal Spray, which will be referred to as Hayfever Relief Spray throughout the rest of this document.

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine is available without prescription. However, you still need to use Hayfever Relief Spray carefully to get the best results from it.
- If any of the side effects gets serious or if you notice any side effects not listed in the leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Hayfever Relief Spray is and what it is used for
2. Before you use Hayfever Relief Spray
3. How to use Hayfever Relief Spray
4. Possible side effects
5. How to store Hayfever Relief Spray
6. Further information

1. WHAT HAYFEVER RELIEF SPRAY IS AND WHAT IT IS USED FOR

Beclometasone Dipropionate is a steroid. It is used for the treatment of seasonal allergic rhinitis (hayfever) in adults aged 18 years of age and over. It acts by reducing the inflammation of your nose which may be the cause of your sneezing, or an itchy, blocked or runny nose.

2. BEFORE YOU USE HAYFEVER RELIEF SPRAY

Do not take Hayfever Relief Spray if you:

- are allergic (hypersensitive) to Beclometasone Dipropionate or any of the other ingredients of Hayfever Relief Spray (see Section 6 for all the ingredients of this medicine).

Do not use if you are under 18 years old.

Take special care and talk to your doctor or pharmacist before using this nasal spray if you:

- are suffering from either a nose infection or sinus trouble
- have recently had surgery or an injury to your nose or ulcers in your nose
- are taking or have recently been treated with other corticosteroid medicines such as asthma inhalers, tablets, injections, nasal sprays, eye or nose drops, creams or ointments.

- are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Some medicines may increase the effects of Hayfever Relief 50 micrograms/dose Nasal Spray and your doctor may wish to monitor you carefully if you are taking these medicines (including some medicines for HIV: ritonavir, cobicistat).

Contact your doctor if you experience blurred vision or other visual disturbances.

If you develop a rash or swelling, stop taking this medicine and tell your doctor immediately.

Pregnancy and breast-feeding

If you are pregnant, think you might be pregnant, intend to become pregnant, or are breast-feeding, speak with your doctor before using this medicine. Ask your doctor or pharmacist for advice before taking any medicine.

Hayfever Relief Spray contains benzalkonium chloride

Benzalkonium chloride may cause irritation or swelling inside the nose, especially if used for a long time.

3. HOW TO USE HAYFEVER RELIEF SPRAY

Always use Hayfever Relief Spray exactly as described in the leaflet.

Using this medicine (Adults aged 18 or over)

Hayfever Relief Spray is not to be used by children or adolescents under 18 years of age.

The recommended dose is two sprays into each nostril twice a day. The maximum daily dose is a total of eight sprays.

Use your nasal spray regularly for maximum relief. It may take a few days for your medicine to work.

Do not exceed the dose stated in this leaflet.

Once your symptoms have improved, you may be able to reduce to one spray into each nostril twice a day. If your symptoms have not improved after 7 days consult your doctor or pharmacist.

Do not use this product continuously for more than 1 month without consulting your doctor.


Method of administration

If you have any problems about using your nasal spray, ask your pharmacist.

1. Remove the dust-cap. Shake the bottle gently.
2. On first using the nasal spray prepare for use by pressing down on the white collar using both your index and middle fingers. Keep the base supported with your thumb (figure 2).

Continue to press down until the collar stops and then allow it to return to its original position. Repeat this action until a fine spray appears. The first five attempts to produce a spray should be allowed to go to waste. Now the spray is ready to use.

3. To use the spray, first blow your nose gently. Closing one nostril off as shown, bend your head forward slightly. Hold the bottle upright and carefully insert the applicator into the other nostril (figure 3).
 4. Slowly begin to breathe in through your nose and whilst doing so press down firmly on the white collar to produce a fine spray inside your nose. Breathe out through your mouth (figure 4).
 5. Repeat step 3 and 4 to squirt a second spray in the same nostril.
- Wash the nozzle frequently with warm water. This will prevent it from getting blocked. If the pump has not been used for a short period of time, re-priming may be necessary (see Step 2 above).



If you use more Hayfever Relief Spray than you should contact your doctor immediately.

If you forget to take Hayfever Relief Spray take a dose as soon as you remember. If it is almost time for your next dose though, do not double the dose, just carry on as before.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Hayfever Relief Spray can cause side effects although not everybody gets them. If you experience one of the following very rare serious side effects stop taking Hayfever Relief Spray and consult your doctor immediately:

- Swelling of your eyes, face, lips or throat
- You develop a rash
- Shortness of breath or wheezing
- You start feeling faint

Other side effects include:

Rare (may affect up to 1 in 1,000 people)
Dryness and irritation of the nose and throat
Unpleasant taste and smell
Bad nose bleeds
Blurred vision
Problems with your eyes such as pain
Damage to your nose

Increased pressure in the eye

A hole or perforation in the nasal septum (the skin tissue which separates the nostrils)

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard
By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE HAYFEVER RELIEF SPRAY

Keep out of the reach and sight of children

Do not use Hayfever Relief Spray after the expiry date which is stated on the label and carton. The expiry refers to the last day of that month.

Discard the contents 3 months after first opening.

- Store below 25°C
- Do not refrigerate
- Protect from light

Medicines should not be disposed of via wastewater or household waste. Ask your Pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

What Hayfever Relief Spray contains

The active substance is Beclometasone Dipropionate and each spray contains 50 micrograms.

The other ingredients are dextrose anhydrous, polysorbate 80, dispersible cellulose, benzalkonium chloride 95%, phenylethanol and purified water.

Hayfever Relief Spray is supplied in a white plastic (high density polyethylene) bottle fitted with a screw on pump covered by a dustcap. The bottle provides 100 sprays.

Marketing Authorisation Holder and Manufacturer

The marketing authorisation is held by Ayrton Saunders Ltd of 9 Arkwright Road, Astmoor Industrial Estate, Runcorn, Cheshire, WA7 1NU.

The medicine is made by Pharmaserve (North West) Ltd, 9 Arkwright Road, Astmoor Industrial Estate, Runcorn, Cheshire. WA7 1NU.

For information in large print, audio CD or braille please contact the Marketing Authorisation Holder.

This leaflet was revised January 2018

LE1773/04