

PRINT LAYOUT DIAGRAM 65MM X 170MM MULTIPEEL WITH 50MM HINGE LEFT

FRONT FACE:



FRONT FACE REVERSE:



- section 3. Liver disease, or severe kidney problems. Shallow breath or breathing difficulties.
 A head injury or persistent headache.
 An asthma attack.
- An astnma attack.

 Previously had a blocked bowel Previously had an addiction to medicines containing Codeine or Morphine If you know that you metabolise very rapidly codeine into morphine Previously had a bad reaction to codeine (a known "ultra-rapid"). Or are under 18 years old • If you are breastfeeding.
- ▲Talk to your doctor or pharmacist if you have.... Kidney or liver problems (a reduced dose may be
- recommended by your doctor) A history of asthma or other breathing problems, including bronchitis Low blood pressure Irritable bowel syndrome (IBS) or other bowel problems
- · Abdominal pain · Myasthenia gravis (a disease of the muscle) . Underactive thyroid or adrenal gland problems (a reduced dose may be recommended by your doctor) • Gallstones or bile duct problems Prostate problems • Problems passing urine
 Irregular heartbeat • Fits • Or you are suffering from shock • Or you are elderly, as elderly people can
- react differently to the medicine. ▲ Talk to your doctor or pharmacist if you are taking....

 • MAOIs (monoamine oxidase inhibitors) for
- depression, selegiline for Parkinson's Disease, or the antibiotic linezolid, or you have taken them within the last two weeks Any tranquillisers, antidepressants, sedatives, painkillers, anaesthetics, sedating antihistamines or sleeping tablets • Any medicine for mental illness • Metoclopramide or domperidone for nausea and vomiting . Quiniding or mexiletine for irregular heartbeat • Alcohol Cimetidine for stomach ulcers • Sodium oxybate for sleeping disorders • Rifampicin for tuberculosis.
- A Pregnant or breastfeeding....
 Codeine Linctus should not be used in pregnancy unless the doctor has told you to do so.
- Do not take codeine while you are breastfeeding. Codeine and morphine passes into breast milk.
- A Driving and using machines...
 This medicine may cause drowsiness or dizziness. If you are affected do not drive or operate machinery. The medicine can affect your ability to drive as it
- may make you sleepy or dizzy.

 Do not drive while taking this medicine until you know how it affects you.

 It is an offence to drive if this medicine affects your ability to drive.
- However, you would not be committing an offence if:
 The medicine has been prescribed to treat a medical or dental problem and
- You have taken it according to the instructions given by the prescriber or in the information provided with the medicine and the way of effective and of the second o olt was not affecting your ability to drive safely. Talk to your doctor or pharmacist if you are not sure whether it is safe for you to drive while taking

92764951905

92764951905

CARRIER:

A Important ingredient information

- This product contains 4g of sucrose per dose. To be taken into account in people with diabetes. It also contains invert syrup. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

 It contains a small amount of ethanol (alcohol), less than 100mg per 5ml.

 It also contains sunset yellow (E110) which may
- cause allergic reaction
- 2. Possible side effects Like all medicines, Codeine Linctus can have side effects, although these don't affect everyone.
- Important side effects
 Codeine is transformed to morphine in the liver by an enzyme. Morphine is the substance that produces pain enzyme. Worpmier's the substance that produces pair relief. Some people have a variation of this enzyme and this can affect people in different ways. In some people, morphine is not produced or produced in very small quantities, and it will not provide enough pain relief. Other people are more likely to get serious side effects because a very high amount of morphine is produced. If you notice any of the following side effects, you must stop taking this medicine and seek immediate medical advice: slow or shallow breathing, confusion.

advice: slow or shallow breathing, confusion,

sleepiness, small pupils, feeling or being sick, constipation, lack of appetite.

- constipation, lack of appetite.

 Other possible side effects are:
 Feeling or being sick Dizziness or vertigo
 Constipation Loss of appetite Fits Feeling unwell
 Muscle twitching or stiffness Drowsiness
 Disturbed sleep Difficulty in passing water
 Dry mouth Sweating Headache Facial flushing
 Fluid retention (e.g. swollen ankles and feet)
 Abnormal heartbeat Feeling faint when standing up
 Slow, shallow breathing Low body heat
 Hallucinations Mood changes Feeling restless or
 confused Pinpoint pupils or other problems with
 eyesight Abdominal pain (including pain caused by
 inflammation of the pancreas (pancreatitis) or bile
 duct problems (biliary spasm)) Ureteric spasm
 (urine tube problem causing abdominal or lower
 back pain) Low sexual urge Allergic reactions
- to addiction, dependence and tolerance (the medicine loses its effect).

 If you notice these or any other side effects, stop use and tell your doctor or pharmacist. They will tell you what to do.

 What the medicine looks like Codeine Linctus is a yellow syrupy liquid. It is supplied in 200ml bottles.

 Text revised: May 2019

this medicine.

- Reporting of side effects
 If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at:

 www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine
- Alf you accidentally take too much see a doctor straight away. Take the pack with you to show which medicine you have swallowed.

 Do not use after the expiry date. The expiry date refers to the last day of that month.

 Return any unused medicine to the pharmacy for disposal The active ingredient per 5ml is: codeine phosphate
- back pain) Low sexual urge Allergic reactions such as itching and a rash Elderly people may experience incontinence, diarrhoea, stomach pain or a blocked bowel Prolonged use may worsen irritable bowel syndrome Prolonged use may lead to addiction, dependence and tolerance (the medicine losse its effect)

 What the medicine looks like Codeina line two is a vellow exprany liquid.

Braille reads:
 c
 o
 d
 e
 i
 n
 e
 l
 i
 n
 c
 t
 u
 s

 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 ••
 <t

PCP Header Box BODY TEXT ONLY

Product Title	Codeine Linctus	Cole	ours Used	Fonts Used	Point Sizes Used Min Max		Proof No.	1	Date		03/06/2019		
Component	Landscape peelable label		Cyan	Helvetica	7pt	9pt							
Pack Size	200ml		Magenta										
IG Code	92764951905		Yellow				CRN	16098	Suffi	x	Α	Approved by	Date
Dimensions	65mm x 170mm		Black				Local Authority	Yes					
Reference Profile Number			Keyline (Non-printing)				submission required						
Barcode	EAN 13 5011309021110		Braille (Non-Printing)				-						
Pharmacode	N/A		Print free area (Non-printing)										
Datamatrix	2764951905		Universal (Non-Printing)								-	Issued by	Date
Verification	2764951905		Cirrus_Info_Box										
Reason for Change	Addition of addiction warning.												