

BACK  
OUTSIDE

FRONT

Please read through the information on the outside and inside of this pack before you start using this medicine. If you have any questions, or if there is anything you do not understand, ask your pharmacist.

**WHAT ARE THEY FOR?**

**Panadol Extra Advance Tablets provide extra effective pain relief:**

- ▼ headache    ▼ migraine    ▼ helps reduce temperature
- ▼ toothache   ▼ sore throat   ▼ aches and pains of colds and flu
- ▼ backache    ▼ period pain   ▼ rheumatic pain   ▼ feverishness

**TAKE SPECIAL CARE**

CONTAINS PARACETAMOL. Do not take with any other paracetamol-containing products.

**Ask your doctor before you take this medicine:**

- if you have liver or kidney disease, including alcoholic liver disease.

**HOW TO TAKE**

Adults and children aged 12 years and over:  
**Swallow 2 tablets every 4 hours as needed.**  
**Do not take more frequently than every 4 hours.**  
**Do not take more than 8 tablets in 24 hours.**  
**Do not give to children under 12 years.**  
**Avoid too much caffeine in drinks like coffee and tea.**  
 High caffeine intake can cause **difficulty sleeping, shaking and an uncomfortable feeling in the chest.**

Do not exceed the stated dose.

Immediate medical advice should be sought in the event of an overdose, even if you feel well, because of the risk of delayed, serious liver damage.

**If symptoms persist consult your doctor.**

**INGREDIENTS**

Each tablet contains the active ingredients Paracetamol 500 mg and Caffeine 65 mg. Also includes a mixture of sodium methyl (E 219), sodium ethyl (E 215) and sodium propyl (E 217) parahydroxybenzoates.

Store below 25°C.

**Keep out of the reach and sight of children.**  
 GlaxoSmithKline Consumer Healthcare (UK) Trading Limited, Brentford, TW8 9GS, U.K.

 5 000347 045219

 GlaxoSmithKline PL 44673/0078 62000000005838

**Panadol**  
*Extra Advance* 500 mg/65 mg Tablets

**Panadol**<sup>®</sup>

**Extra Advance**

500 mg/65 mg Tablets  
Paracetamol + Caffeine

WITH **OPTIZORB** FORMULATION

Fast absorption ✓  
Migraine headache ✓  
Back and muscular aches ✓  
Easy on the stomach ✓

 14

BN: DO NOT USE AFTER THE EXP. DATE:  14

INSIDE

**1. WHAT PANADOL EXTRA ADVANCE TABLETS DO**

**Panadol Extra Advance Tablets are faster absorbed than standard paracetamol and caffeine tablets and are used for the relief of headache, migraine, backache, rheumatic pain, toothache and period pain.** They also relieve sore throat and the fever, aches and pains of colds and flu.

The tablets contain two active ingredients. Paracetamol is a painkiller and also reduces your temperature when you have a fever. Caffeine acts to further help the effectiveness of paracetamol.

**2. CHECK BEFORE YOU TAKE PANADOL EXTRA ADVANCE TABLETS**

**X Do not take Panadol Extra Advance Tablets:**

- if you have ever had an allergic reaction to paracetamol, caffeine or to any of the other ingredients (listed in Section 4).

**! If you are taking other medicines**

Talk to your doctor or pharmacist before taking these tablets if you are taking any prescribed medicines, particularly **metoclopramide** or **domperidone** (for nausea [feeling sick] or vomiting [being sick]) or **colestyramine** (to lower blood cholesterol). If you take **blood thinning drugs** (anticoagulants e.g. **warfarin**) and you need to take a pain reliever on a daily basis, **talk to your doctor** because of the risk of bleeding. But you can still take **occasional** doses of Panadol Extra Advance Tablets at the same time as anticoagulants.

**! Pregnancy and breast feeding**

Due to the caffeine content of this product **it should not be used** if you are **pregnant or breast feeding**.

**3. POSSIBLE SIDE EFFECTS**

Like all medicines Panadol Extra Advance Tablets can have side effects but not everybody gets them. A small number of people have had side effects. **Stop taking** the medicine and

**tell your doctor immediately** if you experience:

- **Allergic reactions** which may be severe such as skin rash and itching sometimes with swelling of the mouth or face or shortness of breath
- **Skin rash or peeling, or mouth ulcers**
- **Breathing problems.** These are more likely if you have experienced them before when taking other painkillers such as ibuprofen and aspirin
- **Unexplained bruising or bleeding**
- **Nausea, sudden weight loss, loss of appetite and yellowing of the eyes and skin.**

**If you do get any side effects**, even those not mentioned in this information, tell your doctor or pharmacist.

If your symptoms continue or your headache becomes persistent, **see your doctor.**

**4. FURTHER INFORMATION**

**Active ingredients:** Each tablet contains Paracetamol 500 mg and Caffeine 65 mg.

**Other ingredients:** Starch pregelatinised, povidone K-25, calcium carbonate, crospovidone, alginate acid, magnesium stearate and sodium methyl (E 219), sodium ethyl (E 215) and sodium propyl (E 217) parahydroxybenzoates.

The tablet coating contains titanium dioxide (E 171), hypromellose, macrogol, polysorbate 80 and carnauba wax.

**The parahydroxybenzoates may cause allergic reactions (possibly delayed).**

Packs of Panadol Extra Advance Tablets contain 8, 14, or 32 tablets.

**The marketing authorisation holder** is GlaxoSmithKline Consumer Healthcare (UK) Trading Limited, Brentford, TW8 9GS, U.K. and all enquiries should be sent to this address.

**The manufacturer** is GlaxoSmithKline Dungarvan Ltd., Co. Waterford, Ireland.

This information was last revised in January 2016.


**Panadol, Optizorb** and the **Triangle device** are registered trademarks of the GSK group of companies.

62000000005838 

TEXT SIZE CONTAINED IN THIS ARTWORK


Body text size: 6.0pt    Smallest text size: 5.0pt    Microtext: N

Dungarvan - Additional Artwork Information Panel





BACK  
OUTSIDE

FRONT



INSIDE



TEXT SIZE CONTAINED IN THIS ARTWORK


Body text size: 6.0pt

Smallest text size: 5.0pt

Microtext: N

BACK  
OUTSIDE

FRONT



INSIDE

**1. WHAT PANADOL EXTRA ADVANCE TABLETS DO**

**Panadol Extra Advance Tablets are faster absorbed than standard paracetamol and caffeine tablets and are used for the relief of headache, migraine, backache, rheumatic pain, toothache and period pain.** They also relieve sore throat and the fever, aches and pains of colds and flu. The tablets contain two active ingredients. Paracetamol is a painkiller and also reduces your temperature when you have a fever. Caffeine acts to further help the effectiveness of paracetamol.

**2. CHECK BEFORE YOU TAKE PANADOL EXTRA ADVANCE TABLETS**

**X Do not take Panadol Extra Advance Tablets:**

- if you have ever had an allergic reaction to paracetamol, caffeine or to any of the other ingredients (listed in Section 4).

**! If you are taking other medicines**  
Talk to your doctor or pharmacist before taking these tablets if you are taking any prescribed medicines, particularly **metoclopramide** or **domperidone** (for **nausea** [feeling sick] or **vomiting** [being sick]) or **colestyramine** (to lower **blood cholesterol**). If you take **blood thinning drugs** (anticoagulants e.g. **warfarin**) and you need to take a pain reliever on a daily basis, **talk to your doctor** because of the risk of bleeding. But you can still take **occasional** doses of Panadol Extra Advance Tablets at the same time as anticoagulants.

**! Pregnancy and breast feeding**  
Due to the caffeine content of this product **it should not be used** if you are **pregnant** or **breast feeding**.

**3. POSSIBLE SIDE EFFECTS**

Like all medicines Panadol Extra Advance Tablets can have side effects but not everybody gets them. A small number of people have had side effects. **Stop taking** the medicine and

**tell your doctor immediately** if you experience:

- **Allergic reactions** which may be severe such as skin rash and itching sometimes with swelling of the mouth or face or shortness of breath
- **Skin rash or peeling**, or **mouth ulcers**
- **Breathing problems**. These are more likely if you have experienced them before when taking other painkillers such as ibuprofen and aspirin
- **Unexplained bruising or bleeding**
- **Nausea**, sudden **weight loss**, **loss of appetite** and **yellowing of the eyes and skin**.

**If you do get any side effects**, even those not mentioned in this information, tell your doctor or pharmacist. If your symptoms continue or your headache becomes persistent, **see your doctor**.

**4. FURTHER INFORMATION**

**Active ingredients:** Each tablet contains Paracetamol 500 mg and Caffeine 65 mg.

**Other ingredients:** Starch pregelatinised, povidone k-25, calcium carbonate, crospovidone, alginate acid, magnesium stearate and sodium methyl (E 219), sodium ethyl (E 215) and sodium propyl (E 217) parahydroxybenzoates. The tablet coating contains titanium dioxide (E 171), hypromellose, macrogol, polysorbate 80 and carnauba wax.

**The parahydroxybenzoates may cause allergic reactions (possibly delayed).**

Packs of Panadol Extra Advance Tablets contain 8, 14, or 32 tablets.


**The marketing authorisation holder** is GlaxoSmithKline Consumer Healthcare (UK) Trading Limited, Brentford, TW8 9GS, U.K. and all enquiries should be sent to this address.

**The manufacturer** is GlaxoSmithKline Dungarvan Ltd., Co. Waterford, Ireland.

This information was last revised in January 2016.

**Panadol**, **Optizorb** and the **Triangle device** are registered trademarks of the GSK group of companies.

62000000005838 glaxoSmithKline



TEXT SIZE CONTAINED IN THIS ARTWORK

Body text size: 6.0pt | Smallest text size: 5.0pt | Microtext: N

Dungarvan - Additional Artwork Information Panel



