

Reference No: TR1307208

Pack Type: Leaflet  
Pack Size: 12/24 Pack  
Action: A

Ref No: PIL 135 x 197

Colours
Process Black
1
LTN

Font Size Panel	
Main Text (body copy):	10 pt
Height (based on lower case x):	1.7 mm

Rennie®

Heartburn, Indigestion & Wind Relief

680 mg/80 mg/25 mg Chewable Tablets

BAYER

Calcium Carbonate, Magnesium Carbonate, Simeticone

Read all of this leaflet carefully because it contains important information for you.

This medicine is available without prescription. However, you still need to take Rennie Heartburn, Indigestion & Wind Relief tablets carefully to get the best results from them. Keep this leaflet, you may need to read it again. Ask your pharmacist if you need more information or advice. You must contact a doctor if your symptoms worsen or do not improve. If you have any unusual effects after taking Rennie Heartburn, Indigestion & Wind Relief, tell your doctor or pharmacist.

IN THIS LEAFLET

1. What is Rennie Heartburn, Indigestion & Wind Relief and what is it used for?
2. Before you take Rennie Heartburn, Indigestion & Wind Relief
3. How to take Rennie Heartburn, Indigestion & Wind Relief
4. Possible side effects
5. How to store Rennie Heartburn, Indigestion & Wind Relief
6. Further information

1. WHAT IS RENNIE® HEARTBURN, INDIGESTION & WIND RELIEF AND WHAT IS IT USED FOR?

Rennie Heartburn, Indigestion & Wind Relief is an antacid tablet which neutralises excess acid to relieve acid indigestion, nervous indigestion, heartburn, flatulence, upset stomach, biliousness; and indigestion and heartburn during pregnancy.

It is also formulated with simeticone, a highly effective ingredient which relieves painful trapped wind and its associated symptoms of uncomfortable bloating, pressure and fullness after food. It acts gently to disperse the trapped air bubbles thus helping to eliminate the symptoms discreetly and without embarrassment.

2. BEFORE YOU TAKE RENNIE® HEARTBURN, INDIGESTION & WIND RELIEF

Do not take Rennie Heartburn, Indigestion & Wind Relief:  
• If you have severe kidney disease.

- If you have high calcium in the urine or kidney stones.
- If you have high calcium or low phosphate levels in the blood.
- If you are allergic (hypersensitive) to any of the ingredients (listed in Section 6. Further Information).

Special precautions:

Do not exceed the stated dose. Rennie Heartburn, Indigestion & Wind Relief should not be used for more than 14 days without consulting a doctor. Please consult your doctor or pharmacist if you have mild or moderate kidney problems as regular blood tests should be carried out. This product contains sorbitol (E420). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

Taking Rennie Heartburn, Indigestion & Wind Relief with food and drink

Avoid taking Rennie Heartburn, Indigestion & Wind Relief with large amounts of milk or dairy products.

Taking other medicines:

- Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.
- If you are taking any of the following medicines speak to your doctor or pharmacist before taking Rennie Heartburn, Indigestion & Wind Relief because it can affect how these medicines work.
  - Tetracyclines and quinolones (antibiotics)
  - Levothyroxine (a thyroid hormone)
  - Eltrombopag (used to increase the number of platelets)
  - Digoxin (used for some heart conditions)
  - Thiazide diuretics (water tablets to treat fluid retention and high blood pressure)
  - Fluorides (for preventing dental caries)
  - Phosphates (for clearing the bowel before a surgical procedure)
  - Iron supplements
- To maximise the benefit of all medicines being used, take Rennie Heartburn, Indigestion & Wind Relief at least 4 hours before or after taking eltrombopag and 1 to 2 hours after taking any other medicines.

3. HOW TO TAKE RENNIE® HEARTBURN, INDIGESTION & WIND RELIEF

Adults and children over 12 years only: 1-2 tablets to be sucked or chewed, up to a maximum of 11 tablets a day.

Children under 12 years: Not recommended.

If symptoms persist after 14 days consult your pharmacist or doctor. Prolonged use should be avoided.

Pregnancy and breast-feeding:

Rennie Heartburn, Indigestion & Wind Relief can be used during pregnancy and breast-feeding if taken as instructed.

If you take more Rennie Heartburn, Indigestion & Wind Relief than you should:

Drink plenty of water and consult your doctor or pharmacist. Symptoms of an overdose include nausea and vomiting, tiredness, increased urine production, increased thirst, dehydration and abnormal muscular weakness.

If you have any questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Rennie Heartburn, Indigestion & Wind Relief can cause side effects, although not everybody gets them. Side effects are unlikely at the recommended dose.

If you experience the following side effects stop treatment immediately and seek medical advice.

Rarely allergic reactions to ingredients have been reported, e.g. rashes, itching, difficulty in breathing and swelling of the face, mouth or throat and anaphylactic shock (anaphylactic shock is a severe sudden allergic reaction, symptoms of which are low blood pressure, shock, palpitations, difficulty in breathing, bronchospasm, skin reactions, abdominal pain or cramps, vomiting and diarrhoea).

Long term use of high doses can cause high blood levels of calcium and magnesium, especially in people with kidney conditions. Symptoms of this may include nausea, vomiting, upset stomach, diarrhoea, tiredness, muscular weakness, headache, kidney problems and impaired sense of taste. In exceptional cases, long term use of high doses can lead to milk alkali syndrome, which can cause high blood levels of calcium.

Reporting of side effects:

If you get any side effects, talk to your doctor,

pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard). By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE RENNIE® HEARTBURN, INDIGESTION & WIND RELIEF

Keep out of the sight and reach of children. Do not store above 25°C and keep in the original pack. Do not use Rennie Heartburn, Indigestion & Wind Relief after the expiry date which is stated on the carton and blister. The expiry date refers to the last day of that month. Medicines should not be disposed of via waste water or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

What Rennie Heartburn, Indigestion & Wind Relief contains:

Each tablet contains the active ingredients: simeticone 25mg, calcium carbonate 680mg and heavy magnesium carbonate 80mg.

The other ingredients are sorbitol (E420), magnesium stearate, potato starch, pregelatinised maize starch, talc, mint flavour, lemon flavour.

What Rennie Heartburn, Indigestion & Wind Relief looks like and the contents of the pack:

Rennie Heartburn, Indigestion & Wind Relief are cream white flat round tablets, scored on one side.

Rennie Heartburn, Indigestion & Wind Relief are available in packs of 2, 4, 6, 12, 16, 18, 20, 24, 36, 48, 60, 96 tablets. Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer:

Marketing Authorisation Holder: Bayer plc, 400 South Oak Way, Reading, RG2 6AD, UK. [consumerhealthuk@bayer.com](mailto:consumerhealthuk@bayer.com)  
Manufacturer: Delpharm Gaillard, 74240 Gaillard, France.

This leaflet was last revised in August 2017