


BELLS	About This Product Bell's Liquid Paraffin BP is taken for the temporary relief of constipation.	Side Effects Prolonged use should be avoided as it may cause anal irritation, inflammation and formation of lumps. Accidentally breathing in liquid paraffin may cause congestion of the lungs. Reporting of side effects: If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this label. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard . By reporting side effects you can help provide more information on the safety of medicines.
	Dosage Liquid for oral use. Adults, the elderly and children over 12 years: Take two to six 5ml spoonfuls (10ml to 30ml). Children from 3 to 12 years: Take one to four 5ml spoonfuls (5ml to 20ml). The above doses may be taken when required. Do not give to children under 3 years.	Active Ingredients Liquid Paraffin 100% What the medicine looks like Liquid Paraffin is a colourless oily liquid. It is supplied in 150ml bottles.
LIQUID PARAFFIN B.P.  For temporary relief of constipation 150ml	Warnings Do not use if you are allergic to Liquid Paraffin. Do not use for long periods. Do not use if you are pregnant or breast-feeding.	
		Further Warnings Consult your doctor or pharmacist if you have persistent abdominal pain or have a condition that makes swallowing difficult. May interfere with the absorption of fat soluble vitamins.
		Storage Keep out of the sight and reach of children. Do not store above 25°C. Protect from light.
		Barcode 5 017848 124942 81 CIN:3800 Ver.1 Revised: Apr. 2016
		Manufacturer and Product Licence Holder: Bell, Sons & Co. (Druggists) Ltd., Southport, PR9 9AL, England. PL3105/0040
		PIP 107-1489
		PEEL BACK AT RED TAB FOR FURTHER IMPORTANT INFORMATION. DO NOT REMOVE.

	Warnings Do not use if you are allergic to Liquid Paraffin. Do not use for long periods. Do not use if you are pregnant or breast-feeding.
	Further Warnings Consult your doctor or pharmacist if you have persistent abdominal pain or have a condition that makes swallowing difficult. May interfere with the absorption of fat soluble vitamins.
	Storage Keep out of the sight and reach of children. Do not store above 25°C. Protect from light.

--	--



BELLS

LIQUID PARAFFIN B.P.



Dispensary Pack
500ml

About This Product

Bell's Liquid Paraffin BP is taken for the temporary relief of constipation.

Dosage

Liquid for oral use.

Adults, the elderly and children over 12 years:

Take two to six 5ml spoonfuls (10ml to 30ml).

Children from 3 to 12 years: Take one to four 5ml spoonfuls (5ml to 20ml).

The above doses may be taken when required.

Do not give to children under 3 years.

Side Effects

Prolonged use should be avoided as it may cause anal irritation, inflammation and formation of lumps. Accidentally breathing in liquid paraffin may cause congestion of the lungs. If you notice these or any other side effects, stop using the product and consult your doctor or pharmacist.

Reporting of side effects:

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this label. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard

By reporting side effects you can help provide more information on the safety of medicines.

Warnings

Do not use if you are allergic to Liquid Paraffin. Do not use if you are pregnant or breast-feeding. Do not use for long periods.

Consult your doctor or pharmacist if you have persistent abdominal pain or have a condition that makes swallowing difficult.

May interfere with the absorption of fat soluble vitamins.

Active Ingredients

Liquid Paraffin 100%

What the medicine looks like

Liquid Paraffin is a colourless oily liquid. It is supplied in 500ml bottles.

Storage

Keep out of the sight and reach of children.

Do not store above 25°C.

Protect from light.

PP 107-1463
PL3105/0040



5 017848 249442
CT15820 Ver. G Revisect-Apr 2016

Manufacturer and
Product Licence
Holder: Bell, Sons & Co.
(Druggists) Ltd.,
Southport,
PR9 9AL, England.