
MHRA Header Box

Product Title	Galenphol Lii	nctus		Co	lours Used	Fonts Used	Point S Min	zes Used Max
i roduct ritie					Process Black	Dax	8.5pt	26pt
Component	Portrait single	e ply labe	el		PANTONE 151 C	Helvetica	6pt	6pt
Pack Size	2 Litre				PANTONE 424 C	OCRB	8.57pt	8.57pt
IG Code	22894810				PANTONE 287 C			
Dimensions	100 x 60mm				Keyline (Non-printing)			
Proof No.	2	Date	11.12.2017					

Datamatrix: 22894810 VERIFIED







FRONT FACE REVERSE:

This medicine contains pholcodine monohydrate, which reduces the desire to cough. It relieves dry coughs which are without phlegm

1. Before you use this medicine 🖾 Do not use the medicine if you have....

• An allergy to any of the ingredients listed in

- section 3.
- Liver disease.
- Shallow breath or breathing difficulties. An asthma attack.
- Taken MAOIs (monoamine oxidase inhibitors)
- for depression within the last two weeks. • Or in children under 12 years old.

A Talk to your doctor or pharmacist if you

- have.... Kidney problems.
- Liver problems.
- A history of asthma or other breathing problems including bronchitis.

92767651901

• A cough for a long time or a cough which produces lots of phlegm. A Talk to your doctor or pharmacist if you

- are taking... Medicine for high blood pressure, e.g. ACE inhibitors, beta blockers.
- Diuretics, which help remove fluid from the body, e.g. furosemide, hydrochlorothiazide.
- · Medicines used to treat sleeplessness, anxiety and schizophrenia. Alcohol or tranquillisers.
- Pain killers, e.g. morphine.

 Or are having surgery, as muscle relaxants used may react with this medicine. Concomitant use of this product and sedative medicines such as benzodiazepines or related drugs increases the risk of drowsiness,

difficulties in breathing (respiratory depression), coma and may be life-threatening. Because of this, concomitant use should only be considered when other treatment options are not possible.

However if your doctor does prescribe this product together with sedative medicines the dose and duration of concomitant treatment should be limited by your doctor. Please tell your doctor about all sedative medicines you are taking, and follow your doctor's dose recommendation closely. It could be helpful to inform friends or relatives to be aware of the signs and symptoms stated above. Contact your doctor when experiencing such symptoms.

A Pregnant or breastfeeding....

Ask your doctor or pharmacist for advice before using this medicine if you are pregnant, might be pregnant or are breastfeeding. Care Pholcodine Oral Solution Sugar Free should not be

used in pregnancy unless the doctor has told you to do so. If you use it when breastfeeding, pholcodine monohydrate in this medicine will be present in small amounts in your milk.

ADriving and using machines....

This medicine may cause drowsiness. If you are affected do not drive or operate machinery.

MHRA Header Box	Вох							
Droduct Title		line 5mg.	Care Pholcodine 5mg/5ml Oral Solution Colours Used	Colc	ours Used	Fonts Used	Point Sizes Used Min Max	es Used Max
	Sugar Free				Process Cyan	Helvetica LT	7pt	9pt
Component	Landscape peelable label	eelable la	abel		Process Magenta			
Pack Size	200ml				Process Yellow			
IG Code	92767651901	_			Process Black			
Dimensions	65mm x 170mm	шш		\bigcirc	Keyline (Non-printing)			
Proof No.	1	Date	31.01.2019	•	Braille			
					Print Free Area (Non-printing)			

CARRIER:

A Important ingredient information

- This medicine contains small amounts of ethanol (alcohol), less than 100mg per 10ml dose.
- It also contains sodium parahydroxybenzoates and amaranth dye which may cause allergic reactions (possibly delayed).

2. Possible side effects

Like all medicines, Care Pholcodine Oral Solution Sugar Free can have side effects, although these don't affect everyone.

Important side effects:

If you think you have any of the following side effects or symptoms, stop using this medicine immediately and see a doctor as soon as possible Allergic reactions which cause difficulty in breathing, fever, swelling of the mouth, lips or skin, severe rashes and itching.

Other possible side effects are: • Feeling sick. Constipation

• Dizziness. Drowsiness · Getting excited. Confusion.

 Being sick. · Rash.

• Failure to cough up phlegm.

If you notice these or any other side effects, stop use and tell your doctor or pharmacist. They will tell you what to do.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for 'MHRA Yellow Card' in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine 3. Further information

A If you accidentally take too much see a doctor straight away. Take the pack with you to show which medicine you have swallowed.

• Do not use after the expiry date. The expiry date refers to the last day of that month.

• Return any unused medicine to the pharmacy

for disposal The active ingredient per 5ml is: pholcodine monohydrate 5mg.

The other ingredients are: citric acid monohydrate (E330), sodium nefyl, ethyl and propyl parahydroxybenzoates (E215, E219, E217), glycerol (E422), ethanol, amaranth dye (E123), saccharin sodium, carmellose sodium, levomenthol, condensed milk flavour, aniseed flavour and purified water.

What the medicine looks like Care Pholcodine Oral Solution Sugar Free is a thick red

liquid which smells of menthol and aniseed. it is supplied in 200ml bottles.

This label was last revised in January 2019 92767651901

